Almost sure limit theorems on Wiener chaos: the non-central case

Ehsan Azmoodeh Ruhr-Universität Bochum, Fakultät für Mathematik ehsan.azmoodeh@rub.de

Ivan Nourdin
Université du Luxembourg, Unité de Recherche en Mathématiques
ivan.nourdin@uni.lu

Abstract

In [1], a framework to prove almost sure central limit theorems for sequences (G_n) belonging to the Wiener space was developed, with a particular emphasis of the case where G_n takes the form of a multiple Wiener-Itô integral with respect to a given isonormal Gaussian process. In the present paper, we complement the study initiated in [1], by considering the more general situation where the sequence (G_n) may not need to converge to a Gaussian distribution. As an application, we prove that partial sums of Hermite polynomials of increments of fractional Brownian motion satisfy an almost sure limit theorem in the long-range dependence case, a problem left open in [1].

Keywords: Almost sure limit theorem; Multiple Wiener-Itô integrals; Malliavin Calculus; Characteristic function; Wiener chaos; Hermite distribution; Fractional Brownian motion.

MSC 2010: 60F05, 60G22, 60G15, 60H05, 60H07.

1 Introduction and main results

Let us start with the following definition, which plays a pivotal role in the paper.

Definition 1.1 Let (G_n) be a sequence of real-valued random variables defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and let μ be a probability measure on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$.

1. We say that (G_n) satisfies an <u>almost sure limit theorem</u> (in short: <u>ASLT</u>) with limit μ if

$$\mathbb{P}\left(\frac{1}{\log n}\sum_{k\leq n}\frac{1}{k}\delta_{G_k}\Rightarrow\mu\ as\ n\to\infty\right)=1,\tag{1.1}$$

where δ_x denotes the Dirac point measure of x and ' \Rightarrow ' stands for the weak convergence of measures.

2. If (1.1) holds true with $\mu = N(0,1)$ (the standard Gaussian distribution), we say that (G_n) satisfies an <u>almost sure central limit theorem</u> (in short: <u>ASCLT</u>).

When μ has a density, it follows from a classical approximation argument that (1.1) is equivalent to saying that

$$\mathbb{P}\left(\lim_{n\to\infty}\frac{1}{\log n}\sum_{k\le n}\frac{1}{k}\mathbf{1}_{\{G_k\leqslant x\}} = \mu(-\infty, x] \text{ for all } x\in\mathbb{R}\right) = 1.$$
 (1.2)

Convergence (1.1) is a kind of weighted strong law of large numbers, but related to a quantity G_n converging in law to μ ; hence its name. Historically, the first appearance of an ASLT was in the book [14] by Lévy. It was stated without proof in the simple situation where $\mu = N(0,1)$ and G_n has the form $G_n = (X_1 + \ldots + X_n)/\sqrt{n}$, for i.i.d. summands X_i with mean 0 and variance 1. Lévy's ASCLT went unnoticed for more than a half century, until it was exploited by Brosamler [7] and Schatte [25] independently in 1998. Since then, there has been a lot of activities around the ASLTs in various contexts, see e.g. [2, 3, 10, 11, 12, 24] and references therein.

We stress on the fact that the validity (or not) of (1.1) is a property of the distribution of the whole sequence (G_n) . Let us elaborate on this point. If $G_n \Rightarrow \mu$, it is well-known by Skorokhod's representation theorem that there exists a family $(G_n^\star)_{n\in\mathbb{N}\cup\{\infty\}}$ of random variables defined on a common probability space such that $G_n^\star \stackrel{\text{law}}{=} G_n$ for all n and $G_n^\star \to G_\infty^\star \sim \mu$ almost surely. Moreover, Cesàro summation theorem implies that, almost surely, $(\log n)^{-1} \sum_{k \leq n} \varphi(G_k^\star)/k \longrightarrow \varphi(G_\infty^\star)$ for any continuous bounded function $\varphi : \mathbb{R} \to \mathbb{R}$; that is, (G_n^\star) does not satisfy an ASLT except if μ is a Dirac mass, since apart in this case $\varphi(G_\infty^\star) \neq \mathbb{E}[\varphi(G_\infty^\star)]$ for at least one function φ .

Now we have made these preliminary comments on general ASLTs, let us concentrate on the specific kind of random variables (G_n) we are interested in in this paper: namely, random variables taking the form of multiple Wiener-Itô integrals. Since the discovery by Nualart and Peccati [21] of their fourth moment theorem (according to which a normalized sequence of multiple Wiener-Itô integrals converges in law to N(0,1) if and only if its fourth moment converges to 3), those stochastic integrals have become a probabilistic object of considerable interest. See for instance the constantly updated webpage [15] for a demonstration of the intense activity surrounding them.

More specifically, a sequence (G_n) of multiple Wiener-Itô integrals converging in law to some μ being given, our goal in this paper is to provide a meaningful set of conditions under which an ASLT holds true. The central case (that is, when $\mu = N(0,1)$) has already been the object of [1]. Therein, a framework has been developed in order to show an ASCLT, by taking advantage of the numerous estimates coming from the Malliavin-Stein approach [18]. One of the main results of [1], that we state here for the sake of comparison, is the following. The notation $I_q(g)$ refers to the qth Wiener-Itô integral with kernel g associated to a given isonormal Gaussian process X over a separable Hilbert space \mathfrak{H} , whereas the notation \otimes_r stands for the contraction operator, see Section 2.3 for more details.

Theorem 1.2 Fix $q \ge 2$, and let (G_n) be a sequence of the form $G_n = I_q(g_n)$, $n \ge 1$. Suppose (for simplicity) that $\mathbb{E}[G_n^2] = 1$ for all n. Suppose in addition that the following two conditions are met.

(A1) for every
$$r \in \{1, \dots, q-1\}$$
: $\sum_{n \geqslant 2} \frac{1}{n \log^2 n} \sum_{k=1}^n \frac{1}{k} \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes(2q-2r)}} < \infty$;

$$(A2) \sum_{n\geqslant 2} \frac{1}{n\log^3 n} \sum_{k,l=1}^n \frac{|\mathbb{E}[G_k G_l]|}{kl} < \infty.$$

Then (G_n) satisfies an ASCLT, that is, (1.1) holds true with $\mu = N(0,1)$.

An application of high interest of Theorem 1.2 studied in [1] concerns the celebrated Breuer-Major theorem. Let $\{X_k\}$ be a stationary centered Gaussian family, characterized by its correlation function $\rho: \mathbb{Z} \to \mathbb{R}$ given by $\rho(k-l) := \mathbb{E}[X_k X_l]$, and assume that

 $\rho(0) = 1$. Fix $q \ge 2$ and let H_q denote the qth Hermite polynomial. Finally, consider the sequence of the Hermite variations of X, defined as

$$V_n = \sum_{k=1}^n H_q(X_k), \quad n \geqslant 1.$$
 (1.3)

It is well known since the seminal works by Breuer and Major [6], Giraitis and Surgailis [9] and Taqqu [27], that the study of the fluctuations of V_n crucially depends on the summability of $|\rho|^q$. More precisely, if we set

$$G_n = \frac{V_n}{\sqrt{\text{Var}(V_n)}}, \quad n \geqslant 1, \tag{1.4}$$

then $G_n \Rightarrow N(0,1)$ as soon as $\sum_{k \in \mathbb{Z}} |\rho(k)|^q < \infty$. When $\sum_{k \in \mathbb{Z}} |\rho(k)|^q = \infty$, the sequence (G_n) may converge to a Gaussian (in some critical situations) or non-Gaussian limit (more likely), see below.

A classical example of a stationary Gaussian sequence $\{X_k\}_{k\in\mathbb{Z}}$ falling within this framework is given by the fractional Gaussian noise

$$X_k = B_{k+1} - B_k, (1.5)$$

where $B = (B_t)_{t \in \mathbb{R}}$ is a fractional Brownian motion of Hurst index $H \in (0,1)$, that is, B is a centered Gaussian process with covariance function

$$\mathbb{E}[B_t B_s] := \frac{1}{2} \left\{ |t|^{2H} + |s|^{2H} - |t - s|^{2H} \right\}, \quad s, t \in \mathbb{R}.$$

More precisely, it is nowadays well-known that

- (i) if $0 < H \le 1 \frac{1}{2q}$ then $G_n \Rightarrow N(0, 1)$;
- (ii) if $1 \frac{1}{2q} < H < 1$ then $G_n \Rightarrow \mu_{H,q}$, the Hermite distribution with parameters H and q.

The proof of (i) follows directly from the Breuer-Major theorem when $H < 1 - \frac{1}{2q}$, whereas a little more effort are required in the critical case $H = 1 - \frac{1}{2q}$ (see, e.g., [5]). For a definition of the Hermite distribution $\mu_{H,q}$ as well as a short proof of the weak convergence (ii) the reader can consult, e.g., [1, Proposition 6.1] and [18, Proposition 7.4.2].

In this paper, we aim to answer the following question: can we associate an almost sure limit theorem to the previous two convergences in law (i) and (ii)? This problem is actually not new, and has been first considered almost one decade ago. In [1], a positive answer has been indeed given for (i), with the help of Theorem 1.2: namely, if $H \leq 1 - \frac{1}{2q}$ and if G_n is defined by (1.4), then (1.1) holds true with $\mu = N(0,1)$. What about the case $H \in (1 - \frac{1}{2q}, 1)$? A solution to this problem was left open in [1] probably because, so far, the Malliavin-Stein approach used therein has been mainly developed in the case of normal approximation.

In the present paper, our main motivation is then to study the case $H \in (1 - \frac{1}{2q}, 1)$ left open in [1]. In order to do so, we first develop a general abstract result (Theorem 1.3), of independent interest and valid in a framework similar to that of Theorem 1.2; then, we apply it to our specific situation (Corollary 1.4).

Theorem 1.3 Fix $q \ge 2$, and let (G_n) be any sequence of the form $G_n = I_q(g_n)$, $n \ge 1$. Suppose $G_n \Rightarrow \mu$, and denote by ϕ_n (resp. ϕ_{∞}) the characteristic function of G_n (resp. μ). Suppose in addition that the following two conditions are met.

(B1) for all
$$r > 0$$
: $\sup_{|t| \le r} \sum_{n \ge 2} \frac{1}{n \log^3 n} \left| \sum_{k=1}^n \frac{1}{k} (\phi_k(t) - \phi_\infty(t)) \right|^2 < \infty$;

(B2)
$$\sum_{n\geqslant 2} \frac{1}{n\log^3 n} \operatorname{Var}\left(\sum_{k=1}^n \frac{G_k^2}{k}\right) < \infty.$$

Then (G_n) satisfies an ASLT, that is, (1.1) holds true.

Let G_n be given by (1.4) with H belonging to $(1 - \frac{1}{2q}, 1)$. In [1, Section 6], the authors have constructed an *explicit* sequence (\widetilde{G}_n) such that $\widetilde{G}_n \stackrel{\text{law}}{=} G_n$ for any n and \widetilde{G}_n does not satisfy an ASLT. But as we already explained in the second paragraph following Definition 1.1, from this there is nothing to learn about the validity or not of an ASLT associated to the original sequence (G_n) . Here, using our abstract Theorem 1.3, we are actually able to prove that (G_n) does satisfy an ASLT.

Corollary 1.4 Let G_n be given by (1.4), with X_k as in (1.5) and $H \in (1 - \frac{1}{2q}, 1)$. Then (G_n) satisfies an ASLT with $\mu = \mu_{H,q}$ the Hermite distribution with parameters H and q.

To conclude this introduction, we would like to stress that our Theorem 1.3 is a true extension of Theorem 1.2, in the sense that the latter can be obtained as a particular case of the former: see Section 5 for the details.

Also, for the sake of illustration, we develop in Section 4 an easy example of application of our Theorem 1.3 involving a sequence of independent standard Gaussian random variables.

The rest of the paper is organized as follows. In Section 2 several preliminary results that are needed for the proofs are collected. Section 3 is devoted to the proof of Theorem 1.3. In Section 4, an easy application of Theorem 1.3 to partial sums of independent standard Gaussian random variables is presented. The proof of Corollary 1.4 can be found in Section 5. Finally, Section 6 details how to deduce [1, Theorem 1.2] from our Theorem 1.3.

2 Preliminaries

2.1 Ibragimov-Lifshits criterion for ASLT

As we will see, we are not going to prove *directly* that (G_n) in Theorem 1.3 satisfies (1.2). Instead, we are going to check the validity of a powerful criterion due to Ibragimov and Lifshits [12], that we recall here for the convenience of the reader.

Theorem 2.1 Let (G_n) be a sequence of real-valued random variables defined on a common probability space and converging in distribution towards μ , and set

$$\Delta_n(t) = \frac{1}{\log n} \sum_{k=1}^n \frac{1}{k} \left(e^{itG_k} - \int_{\mathbb{R}} e^{itx} \mu(dx) \right). \tag{2.6}$$

If

$$\sup_{|t| \le r} \sum_{n} \frac{\mathbb{E}|\Delta_n(t)|^2}{n \log n} < \infty \quad \text{for all } r > 0,$$
 (2.7)

then (G_n) satisfies the ASLT (1.1).

2.2 An easy reduction lemma

In this section, we state and prove an easy reduction lemma, that we are going to use in the proof of Corollary 1.4.

Lemma 2.2 Suppose that (G_n) is such that $G_n \Rightarrow \mu$, and assume that μ has a density. Let (a_n) be a real-valued sequence converging to $a_\infty \neq 0$. Then (G_n) satisfies an ASLT if and only if (a_nG_n) does.

Proof. Without loss of generality, we may and will assume that $a_n > 0$ for all n and that $a_{\infty} = 1$. By symmetry, only the implication "if (G_n) satisfies an ASLT then $(a_n G_n)$ does" has to be proved.

So, let us assume that (G_n) satisfies an ASLT. Since μ has a density, we are going to use criterion (1.2). Fix $\varepsilon > 0$, and let k_0 be such that $1 - \varepsilon \leqslant a_k \leqslant 1 + \varepsilon$ for all $k \geqslant k_0$. For any $x \in \mathbb{R}$, we can write, for all $k \geqslant k_0$,

$$\begin{array}{lll} \mathbf{1}_{\{G_k\leqslant \frac{x}{1+\varepsilon}\}}\mathbf{1}_{\{x\geqslant 0\}} + \mathbf{1}_{\{G_k\leqslant \frac{x}{1-\varepsilon}\}}\mathbf{1}_{\{x< 0\}} & \leqslant & \mathbf{1}_{\{a_kG_k\leqslant x\}} \\ & \leqslant & \mathbf{1}_{\{G_k\leqslant \frac{x}{1-\varepsilon}\}}\mathbf{1}_{\{x\geqslant 0\}} + \mathbf{1}_{\{G_k\leqslant \frac{x}{1+\varepsilon}\}}\mathbf{1}_{\{x< 0\}}. \end{array}$$

As a consequence,

$$\frac{1}{\log n} \sum_{k=k_0}^n \frac{1}{k} \left(\mathbf{1}_{\{G_k \leqslant \frac{x}{1+\varepsilon}\}} \mathbf{1}_{\{x \geqslant 0\}} + \mathbf{1}_{\{G_k \leqslant \frac{x}{1-\varepsilon}\}} \mathbf{1}_{\{x < 0\}} \right) \leqslant \frac{1}{\log n} \sum_{k=1}^n \frac{1}{k} \mathbf{1}_{\{a_k G_k \leqslant x\}}$$

and

$$\frac{1}{\log n} \sum_{k=1}^{n} \frac{1}{k} \mathbf{1}_{\{a_k G_k \leqslant x\}} \leqslant \frac{k_0}{\log n} + \frac{1}{\log n} \sum_{k=k_0}^{n} \frac{1}{k} \left(\mathbf{1}_{\{G_k \leqslant \frac{x}{1-\varepsilon}\}} \mathbf{1}_{\{x \geqslant 0\}} + \mathbf{1}_{\{G_k \leqslant \frac{x}{1+\varepsilon}\}} \mathbf{1}_{\{x < 0\}} \right).$$

Letting $n \to \infty$, we deduce from (1.2) that, almost surely, for all $x \in \mathbb{R}$,

$$\mu(-\infty, (1+\varepsilon)^{-1}x] \, \mathbf{1}_{\{x \ge 0\}} + \mu(-\infty, (1-\varepsilon)^{-1}x] \, \mathbf{1}_{\{x < 0\}} \le \liminf_{n \to \infty} \frac{1}{\log n} \sum_{k=1}^{n} \frac{1}{k} \mathbf{1}_{\{a_k G_k \le x\}}$$

$$\le \limsup_{n \to \infty} \frac{1}{\log n} \sum_{k=1}^{n} \frac{1}{k} \mathbf{1}_{\{a_k G_k \le x\}} \le \mu(-\infty, (1-\varepsilon)^{-1}x] \, \mathbf{1}_{\{x \ge 0\}} + \mu(-\infty, (1+\varepsilon)^{-1}x] \, \mathbf{1}_{\{x < 0\}}.$$

Letting now $\varepsilon \to 0$ yields $\frac{1}{\log n} \sum_{k=1}^n \frac{1}{k} \mathbf{1}_{\{a_k G_k \leq x\}} \to \mu(-\infty, x]$, which is desired conclusion.

2.3 Elements of Malliavin calculus

Our proofs of Theorem 1.3 and Corollary 1.4 are based on the use of the Malliavin calculus. This is why in this section we recall the few elements of Gaussian analysis and Malliavin calculus that will be needed. For more details or missing proofs, we invite the reader to consult one of the three books [18, 22, 23].

Let \mathfrak{H} be a real separable Hilbert space. For any $q \geq 1$, we write $\mathfrak{H}^{\otimes q}$ and $\mathfrak{H}^{\odot q}$ to indicate, respectively, the qth tensor power and the qth symmetric tensor power of \mathfrak{H} .

We denote by $X = \{X(h) : h \in \mathfrak{H}\}$ an isonormal Gaussian process over \mathfrak{H} , that is, X is a centered Gaussian family defined a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$ satisfying $\mathbb{E}[X(g)X(h)] = \langle g, h \rangle_{\mathfrak{H}}$. We also assume that \mathcal{F} is generated by X, and use the shorthand notation $L^2(\Omega) = L^2(\Omega, \mathcal{F}, \mathbb{P})$.

For every $q \ge 1$, the qth Wiener chaos of X is defined as the closed linear subspace of $L^2(\Omega)$ generated by the family $\{H_q(X(h)): h \in \mathfrak{H}, \|h\|_{\mathfrak{H}} = 1\}$, where H_q is the qth Hermite polynomial:

$$H_q(x) = (-1)^q e^{\frac{x^2}{2}} \frac{d^q}{dx^q} \left(e^{-\frac{x^2}{2}} \right). \tag{2.8}$$

For any $q\geqslant 1$, the mapping $I_q(h^{\otimes q})=H_q(W(h))$ (for $h\in\mathfrak{H},\|h\|_{\mathfrak{H}}=1$) can be extended to a linear isometry between the symmetric tensor product $\mathfrak{H}^{\odot q}$ (equipped with the modified norm $\sqrt{q!} \|\cdot\|_{\mathfrak{H}^{\otimes q}}$) and the qth Wiener chaos.

Let $\{e_k, k \geqslant 1\}$ be a complete orthonormal system in \mathfrak{H} . Given $f \in \mathfrak{H}^{\odot p}$ and $g \in \mathfrak{H}^{\odot q}$, for every $r = 0, \dots, p \land q$, the contraction of f and g of order r is the element of $\mathfrak{H}^{\otimes (p+q-2r)}$

$$f \otimes_r g = \sum_{i_1, \dots, i_r = 1}^{\infty} \langle f, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}} \otimes \langle g, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}}.$$
 (2.9)

Notice that the definition of $f \otimes_r g$ does not depend on the particular choice of $\{e_k, k \geq 1\}$, and that $f \otimes_r g$ is not necessarily symmetric; we denote its symmetrization by $f \otimes_r g \in$ $\mathfrak{H}^{\odot(p+q-2r)}$. Moreover, $f\otimes_0 g=f\otimes g$ equals the tensor product of f and g while, for $p=q,\ f\otimes_q g=\langle f,g\rangle_{\mathfrak{H}^{\otimes q}}.$ Contractions appear naturally in the product formula for multiple integrals: if $f \in \mathfrak{H}^{\odot p}$ and $g \in \mathfrak{H}^{\odot q}$, then

$$I_p(f)I_q(g) = \sum_{r=0}^{p \wedge q} r! \binom{p}{r} \binom{q}{r} I_{p+q-2r}(f\widetilde{\otimes}_r g). \tag{2.10}$$

Proof of Theorem 1.3 3

As we said in Section 2.1, we are going to check the condition (2.7) in our specific framework. For $t \in \mathbb{R}$, recall the quantity $\Delta_n(t)$ from (2.6) and that ϕ_k (resp. ϕ_{∞}) stands for the characteristic function of G_k (resp. μ). We also denote by $\varphi_{k,l}$ the characteristic function of $G_k - G_l$. With these notation in mind, we can now write

$$\mathbb{E}|\Delta_{n}(t)|^{2} = \frac{1}{\log^{2} n} \sum_{k,l=1}^{n} \frac{1}{kl} \mathbb{E}\left(\left(e^{itG_{k}} - \phi_{\infty}(t)\right) \times \left(e^{-itG_{l}} - \phi_{\infty}(-t)\right)\right)
= \frac{1}{\log^{2} n} \sum_{k,l=1}^{n} \frac{1}{kl} \left\{\left(\phi_{k}(t) - \phi_{\infty}(t)\right) \left(\phi_{l}(-t) - \phi_{\infty}(-t)\right) + \left(\varphi_{k,l}(t) - \phi_{k}(t)\phi_{l}(-t)\right)\right\}
=: A_{1}(n,t) + A_{2}(n,t).$$

Now, note that

$$A_1(n,t) = \frac{1}{\log^2 n} \left| \sum_{k=1}^n \frac{1}{k} (\phi_k(t) - \phi_\infty(t)) \right|^2,$$

so that $\sup_{|t| \leq r} \sum_{n \geqslant 2} \frac{A_1(n,t)}{n \log n} < \infty$ for all r > 0, by condition (B1) in Theorem 1.3. On the other hand, [16, Proposition 3.1] provides the existence of a universal constant

c such that

$$\left| \varphi_{k,l}(t) - \phi_k(t)\phi_l(-t) \right| \leqslant c|t| \operatorname{Cov}(G_k^2, G_l^2).$$

We deduce that

$$A_2(n,t) \le \frac{c|t|}{\log^2 n} \sum_{k,l=1}^n \frac{1}{kl} \mathbf{Cov}(G_k^2, G_l^2) = \frac{c|t|}{\log^2 n} \mathbf{Var}\left(\sum_{k=1}^n \frac{G_k^2}{k}\right),\tag{3.11}$$

implying in turn by condition (B2) that $\sup_{|t| \leq r} \sum_{n \geq 2} \frac{A_2(n,t)}{n \log n} < \infty$ for all r > 0, and concluding the proof.

4 An easy illustrating example

Before proceeding with the proof of Corollary 1.4, let us illustrate the use of our Theorem 1.3 on a simple example.

Let $X_0, X_1, X_2 \dots \sim N(0, 1)$ be a sequence of independent copies defined on the same probability space, and consider

$$G_n = \frac{X_n}{\sqrt{2n}} \sum_{j=0}^{n-1} (X_j^2 - 1).$$

A first observation is that G_n can be realized as a Wiener-Itô integral of order 3 with respect to an isonormal Gaussian process X over the Hilbert space $\mathfrak{H} = L^2(\mathbb{R}_+)$. Indeed, if we identify X_j with $X(\mathbf{1}_{[j,j+1]})$, we get

$$G_n = I_3(g_n)$$
, where $g_n = \frac{1}{\sqrt{2n}} \sum_{j=0}^{n-1} \widetilde{\mathbf{1}}_{[n,n+1] \times [j,j+1]^2}$,

where the tilde means that the function has been symmetrized.

Thus, we are in a position to make use of Theorem 1.3, with μ the distribution of the product of two independent standard Gaussian random variables.

First, the fact that $G_n \Rightarrow \mu$ is obvious by the classical central limit theorem, since

$$G_n \stackrel{\text{law}}{=} \frac{X_0}{\sqrt{2n}} \sum_{j=1}^n (X_j^2 - 1).$$

It is moreover clear that, for any (fixed) r > 0

$$\sup_{|t| \leq r} \left| \mathbb{E}[e^{itG_k}] - \int_{\mathbb{R}} e^{itx} d\mu \right| = \sup_{|t| \leq r} \left| \mathbb{E}\left[e^{-\frac{t^2}{2}\left\{\frac{1}{\sqrt{2k}}\sum_{j=1}^n (X_j^2 - 1)\right\}^2\right]} - \mathbb{E}\left[e^{-\frac{t^2}{2}X_0^2}\right] \right|.$$

The right-hand side of the previous equality being a $O(\frac{1}{k})$ by the classical Berry-Esseen theorem (e.g. with the Wasserstein distance), the condition (B1) of Theorem 1.3 is immediately satisfied.

We now turn to (B2). Using the product formula (2.10) for multiple integrals, we can write

$$\sum_{k=1}^{n} \frac{G_k^2 - \mathbb{E}[G_k^2]}{k} = I_6 \left(\sum_{k=1}^{n} \frac{1}{k} g_k \widetilde{\otimes} g_k \right) + 9 I_4 \left(\sum_{k=1}^{n} \frac{1}{k} g_k \widetilde{\otimes}_1 g_k \right) + 18 I_2 \left(\sum_{k=1}^{n} \frac{1}{k} g_k \widetilde{\otimes}_2 g_k \right).$$

We are thus left to check that, for $r = \{0, 1, 2\}$:

$$\sum_{n\geqslant 2} \frac{1}{n\log^3 n} \left\| \sum_{k=1}^n \frac{1}{k} g_k \otimes_r g_k \right\|_{\mathfrak{H}^{\otimes(2q-2r)}}^2 < \infty.$$

This is actually immediate, by using the following straightforward computations (left to the reader):

$$\langle g_k \otimes g_k, g_l \otimes g_l \rangle_{\mathfrak{H}^{\otimes 6}} = \frac{(k \wedge l)^2}{36 \, kl}, \quad \langle g_k \otimes_1 g_k, g_l \otimes_1 g_l \rangle_{\mathfrak{H}^{\otimes 4}} = \frac{4 \, k \wedge l + (k \wedge l)^2}{324 \, kl}$$
$$\langle g_k \otimes_2 g_k, g_l \otimes_2 g_l \rangle_{\mathfrak{H}^{\otimes 2}} = \frac{5 \, k \wedge l}{324 \, kl}. \tag{4.12}$$

Summarizing, we have shown the following result.

Proposition 4.1 Let $X_0, X_1, X_2 ... \sim N(0,1)$ be a sequence of independent copies defined on the same probability space, and consider $G_n = \frac{X_n}{\sqrt{2n}} \sum_{j=0}^{n-1} (X_j^2 - 1)$. Then (G_n) satisfies an ASLT with μ the distribution of X_0X_1 .

Remark. It is interesting to observe that the expressions (4.12) are no longer valid if we replace G_n by $\widetilde{G}_n = \frac{X_0}{\sqrt{2n}} \sum_{j=1}^n (X_j^2 - 1)$. Actually, the assumption (B2) of Theorem 1.3 turns out to be not satisfied for (\widetilde{G}_n) . We believe that it is because \widetilde{G}_n does not satisfy an ASLT, see also the discussion in the second paragraph following Definition 1.1.

5 Proof of Corollary 1.4

Let (G_n) be given by (1.4) with $H \in (1 - \frac{1}{2q}, 1)$ and recall V_n from (1.3). It is a straightforward exercise to prove that $\operatorname{Var}(V_n)$ is equivalent to a constant times $n^{2-2q(1-H)}$ as $n \to \infty$ (see, e.g. [1] and references therein). Thanks to Lemma 2.2 and because the Hermite distribution $\mu_{H,q}$ admits a density (this latter fact is an immediate consequence of the main result of [19] for instance), it is then equivalent to prove an ASLT for (G_n) or for (\widehat{G}_n) defined as

$$\widehat{G}_n = n^{q(1-H)-1} \sum_{j=1}^n H_q(B_{j+1} - B_j).$$

In the sequel, we are thus rather going to prove that (\widehat{G}_n) satisfies an ASLT.

Our first observation is that \widehat{G}_n takes the form of a multiple Wiener-Itô integral with respect to B, after identifying this latter with an isonormal Gaussian process X over the Hilbert space \mathfrak{H} defined as the closure of the set of step functions with respect to the scalar product $\langle \mathbf{1}_{[0,t]}, \mathbf{1}_{[0,s]} \rangle_{\mathfrak{H}} = \mathbb{E}[B_t B_s], s, t \geqslant 0$, see [18, Proposition 7.2.3] for further details. More precisely, we have

$$\widehat{G}_n = I_q(g_n)$$
, where $g_n = n^{q(1-H)-1} \sum_{j=1}^n \mathbf{1}_{[j,j+1]}^{\otimes q}$,

This observation being made, we can now use Theorem 1.3, and check that conditions (B1) and (B2) therein are in order. In what follows, ϕ_k and ϕ_{∞} denote the characteristic functions of \hat{G}_k and $\mu_{H,q}$ respectively.

For (B1), it is a direct consequence of some estimates given in [5]. More precisely, using [5, Proposition 3.1] one infers that, for any (fixed) r > 0,

$$\sup_{|t| \leqslant r} |\phi_k(t) - \phi_{\infty}(t)| \leqslant r d_W(\widehat{G}_k, \mu_{H,q}) = O(k^{q(1-H)-\frac{1}{2}}), \tag{5.13}$$

where d_W stands for the Wasserstein distance. The first inequality in (5.13) is a direct consequence of the basic fact that $x \mapsto e^{itx}$ is a Lipschitz-continuous function with Lipschitz constant |t|. Also, recall for a real-valued random variables $F_1 \sim \mu_1$ and $F_2 \sim \mu_2$ that

$$d_W(F_1, F_2) = d_W(\mu_1, \mu_2) = \sup \left\{ \left| \int_{\mathbb{R}} h d\mu_1 - \int_{\mathbb{R}} h d\mu_2 \right| : h : \mathbb{R} \to \mathbb{R} \text{ s.t. } \|h\|_{\text{Lip}} \le 1 \right\}.$$

Therefore, for some constant C, we obtain

$$\sup_{|t| \leqslant r} \sum_{n \geqslant 2} \frac{1}{n \log^3 n} \left| \sum_{k=1}^n \frac{1}{k} (\phi_k(t) - \phi_\infty(t)) \right|^2 \leqslant C \sum_{n \geqslant 2} \frac{1}{n \log^3 n} \left(\sum_{k=1}^n k^{q(1-H) - \frac{3}{2}} \right)^2 < \infty.$$

Let us now turn to (B2). Using the product formula (2.10) for multiple integrals, we can write

$$\sum_{k=1}^{n} \frac{G_k^2}{k} = \sum_{r=0}^{q} r! \binom{q}{r}^2 I_{2q-2r} \left(\sum_{k=1}^{n} \frac{1}{k} g_k \widetilde{\otimes}_r g_k \right).$$

Thus we are left to check that, for all $r \in \{0, \ldots, q-1\}$:

$$\sum_{n \geqslant 2} \frac{1}{n \log^3 n} \left\| \sum_{k=1}^n \frac{1}{k} g_k \otimes_r g_k \right\|_{\mathfrak{H}^{\otimes(2q-2r)}}^2 < \infty.$$
 (5.14)

Let us first focus on the case where $1 \leqslant r \leqslant q-1$. We can write, with $\rho(a) = \frac{1}{2} \left(|a+1|^{2H} + |a-1|^{2H} - 2|a|^{2H} \right)$ and $\rho_k(a) = |\rho(a)| \mathbf{1}_{|a| \leqslant k}$:

$$\left\| \sum_{k=1}^{n} \frac{1}{k} g_{k} \otimes_{r} g_{k} \right\|_{\mathfrak{H}^{\otimes}(2q-2r)}^{2} = \sum_{k,l=1}^{n} \frac{1}{kl} \langle g_{k} \otimes_{r} g_{k}, g_{l} \otimes_{r} g_{l} \rangle_{\mathfrak{H}^{\otimes}(2q-2r)}$$

$$\leq 2 \sum_{1 \leqslant l \leqslant k \leqslant n} (kl)^{2q(1-H)-3} \sum_{i,j=1}^{k} \sum_{i',j'=1}^{l} \rho(i-j)^{r} \rho(i-i')^{q-r} \rho(i'-j')^{r} \rho(j-j')^{q-r}$$

$$\leq 2 \sum_{1 \leqslant l \leqslant k \leqslant n} (kl)^{2q(1-H)-3} \sum_{i=1}^{k} \sum_{j'=1}^{l} (\rho_{k}^{r} * \rho_{k}^{q-r})(i-j')^{2}$$

$$\leq 2 \sum_{1 \leqslant l \leqslant k \leqslant n} (kl)^{2q(1-H)-3} l \|\rho_{k}^{r} * \rho_{k}^{q-r}\|_{\ell^{2}(\mathbb{Z})}^{2}.$$

Using the Young inequality with $p = \frac{2q}{3r}$ and $p' = \frac{2q}{3(q-r)}$ (so that $\frac{1}{p} + \frac{1}{p'} = \frac{3}{2}$), we obtain

$$\|\rho_k^r * \rho_k^{q-r}\|_{\ell^2(\mathbb{Z})}^2 \leqslant \|\rho_k^r\|_{\ell^p(\mathbb{Z})}^2 \|\rho_k^{q-r}\|_{\ell^{p'}(\mathbb{Z})}^2 = \left(\sum_{|j| \leqslant k} |\rho(j)|^{\frac{2q}{3}}\right)^3.$$

But $\rho(j) \sim H(2H-1)|j|^{2H-2}$ and, because $H > 1 - \frac{1}{2q} > 1 - \frac{3}{4q}$, we deduce that

$$\left(\sum_{|j| \le k} |\rho(j)|^{\frac{2q}{3}}\right)^3 = O(k^{3-4q(1-H)}) \text{ as } k \to \infty.$$

Thus, for $r \in \{1, ..., q - 1\}$,

$$\left\| \sum_{k=1}^{n} \frac{1}{k} g_k \otimes_r g_k \right\|_{\mathfrak{H}^{\otimes(2q-2r)}}^2 = O\left(\sum_{1 \leqslant l \leqslant k \leqslant n} l^{2q(1-H)-2} k^{-2q(1-H)} \right) = O\left(\sum_{k=1}^{n} \frac{1}{k} \right)$$

$$= O(\log n). \tag{5.15}$$

Actually, the previous estimate (5.15) is also valid for r=0. Indeed, we have in this case

$$\left\| \sum_{k=1}^{n} \frac{1}{k} g_{k} \otimes_{0} g_{k} \right\|_{\mathfrak{H}^{3}(2q)}^{2} = \sum_{k,l=1}^{n} (kl)^{2q(1-H)-3} \left(\sum_{i=1}^{k} \sum_{j=1}^{l} \rho(i-j)^{q} \right)^{2}$$

$$\leq 2 \sum_{1 \leq l \leq k \leq n} (kl)^{2q(1-H)-3} \left(\sum_{i=1}^{k} \sum_{j=1}^{l} |\rho(i-j)|^{q} \right)^{2}$$

$$\leq 2 \sum_{1 \leq l \leq k \leq n} (kl)^{2q(1-H)-3} l^{2} \left(\sum_{|r| < k} |\rho(r)|^{q} \right)^{2}$$

$$\leq \operatorname{cst} \sum_{1 \leq l \leq k \leq n} (kl)^{2q(1-H)-3} l^{2} k^{2-4(1-H)q} = O(\sum_{k=1}^{n} \frac{1}{k}) = O(\log n), \tag{5.16}$$

where the last inequality used the fact that $\rho(r) \sim H(2H-1)r^{2H-2}$ as $|r| \to \infty$. Finally, (5.15) and (5.16) together imply that (5.14) takes place for any $r \in \{0, \ldots, q-1\}$, which concludes the proof of Theorem 1.4.

6 Theorem 1.3 implies Theorem 1.2

To conclude this paper, let us explain how to deduce Theorem 1.2 (taken from [1]) from our Theorem 1.3, even if at first glance conditions (A1)-(A2) and (B1)-(B2) do not seem to be related to each others.

Fix $q \ge 2$, and let (G_n) be a sequence of the form $G_n = I_q(g_n)$, $n \ge 1$. Suppose moreover that $\mathbb{E}[G_n^2] = q! \|g_n\|_{\mathfrak{H}^{\otimes q}}^2 = 1$ for all n. In what follows, $c_q > 0$ denotes a constant only depending on q, whose value can change from line to line.

Firstly, we deduce from [18, Theorem 5.2.6] (to get the second inequality) and [18, (5.2.6)] (to get the third inequality) that, for all $t \in \mathbb{R}$,

$$\begin{split} \left| \mathbb{E}[e^{itG_k}] - e^{-t^2/2} \right| & \leqslant |t| \, d_W(G_k, N(0, 1)) \leqslant c_q \, |t| \, (\mathbb{E}[G_k^4] - 3) \\ & \leqslant c_q \, |t| \max_{1 \leqslant r \leqslant q - 1} \|g_k \widetilde{\otimes}_r g_k\|_{\mathfrak{H}^{\otimes (2q - 2r)}} \\ & \leqslant c_q \, |t| \max_{1 \leqslant r \leqslant q - 1} \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes (2q - 2r)}}. \end{split}$$

As a consequence,

$$\begin{split} \sup_{|t|\leqslant r} \sum_{n\geqslant 2} \frac{1}{n\log^3 n} \left| \sum_{k=1}^n \frac{1}{k} (\mathbb{E}[e^{itG_k}] - e^{-t^2/2}) \right|^2 \\ \leqslant & 2 \sup_{|t|\leqslant r} \sum_{n\geqslant 2} \frac{1}{n\log^2 n} \left| \sum_{k=1}^n \frac{1}{k} (\mathbb{E}[e^{itG_k}] - e^{-t^2/2}) \right| \\ \leqslant & 2 r c_q \max_{1\leqslant r\leqslant q-1} \sum_{n\geqslant 2} \frac{1}{n\log^2 n} \sum_{k=1}^n \frac{1}{k} \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes(2q-2r)}}, \end{split}$$

from which it comes that (A1) implies (B1).

Secondly, one can write

$$\begin{aligned} \|g_k \otimes_r g_l\|_{\mathfrak{H}^{\otimes}(2q-2r)}^2 &= \langle g_k \otimes_r g_k, g_l \otimes_r g_l \rangle_{\mathfrak{H}^{\otimes}(2q-2r)} \\ &\leqslant \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes}(2r)} \|g_l \otimes_r g_l\|_{\mathfrak{H}^{\otimes}(2q-2r)} \\ &\leqslant \|g_l\|_{\mathfrak{H}^{\otimes}(2q-2r)}^2 = q! \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes}(2q-2r)}. \end{aligned}$$

Moreover, it has been shown in [20, Proof of Theorem 4.3] that

$$\mathbf{Cov}(G_k^2, G_l^2) - 2 \left(\mathbb{E}[G_k G_l] \right)^2$$

$$= q!^2 \sum_{r=1}^{q-1} {q \choose r}^2 \|g_k \otimes_r g_l\|_{\mathfrak{H}^{\otimes}(2q-2r)}^2 + \sum_{r=1}^{q-1} r!^2 {q \choose r}^4 (2q-2r)! \|g_k \widetilde{\otimes}_r g_l\|_{\mathfrak{H}^{\otimes}(2q-2r)}^2.$$

We deduce that

$$0 \leqslant \mathbf{Cov}(G_k^2, G_l^2) \leqslant 2 \left(\mathbb{E}[G_k G_l] \right)^2 + c_q \max_{1 \leqslant r \leqslant q-1} \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes(2q-2r)}}$$
$$\leqslant 2 \left| \mathbb{E}[G_k G_l] \right| + c_q \max_{1 \leqslant r \leqslant q-1} \|g_k \otimes_r g_k\|_{\mathfrak{H}^{\otimes(2q-2r)}},$$

implying in turn that

$$\sum_{n\geqslant 2} \frac{1}{n \log^{3} n} \operatorname{Var}\left(\sum_{k=1}^{n} \frac{G_{k}^{2}}{k}\right) = \sum_{n\geqslant 2} \frac{1}{n \log^{3} n} \sum_{k,l=1}^{n} \frac{1}{k l} \operatorname{Cov}\left(G_{k}^{2}, G_{l}^{2}\right)$$

$$\leqslant 2 \sum_{n\geqslant 2} \frac{1}{n \log^{3} n} \sum_{k,l=1}^{n} \frac{|\mathbb{E}[G_{k}G_{l}]|}{k l} + c_{q} \max_{1\leqslant r\leqslant q-1} \sum_{n\geqslant 2} \frac{1}{n \log^{2} n} \sum_{k=1}^{n} \frac{1}{k} \|g_{k} \otimes_{r} g_{k}\|_{\mathfrak{H}^{\otimes}(2q-2r)},$$

from which it follows that (A1) and (A2) imply (B2).

Acknowledgment. We thank two anonymous referees for their useful comments that led to an improvement in the presentation of this paper.

References

- [1] B. Bercu, I. Nourdin and M. Taqqu (2010): Almost sure central limit theorems on the Wiener space. *Stoch. Proc. Appl.* **120**, pp. 1607-1628.
- [2] I. Berkes and E. Csáki (2001): A universal result in almost sure central limit theory. Stoch. Process. Appl. 94, no. 1, pp. 105-134.
- [3] I. Berkes and H. Dehling (1993): Some limit theorems in log density. Ann. Probab. 21, pp. 1640-1670.
- [4] I. Berkes and L. Horváth (1999): Limit theorems for logarithmic averages of fractional Brownian motions. J. Theoret. Probab. 12, no. 4, pp. 985-1009.
- [5] J.-C. Breton and I. Nourdin (2008): Error bounds on the non-normal approximation of Hermite variations of fractional Brownian motion. *Electron. Comm. Probab.* **13**, pp. 482-493.
- [6] P. Breuer and P. Major (1983): Central limit theorems for nonlinear functionals of Gaussian fields, *J. Multivariate Anal.* **13**, no. 3, pp. 425-441.
- [7] G.A. Brosamler (1988): An almost everywhere central limit theorem. *Math. Proc. Cambridge Philos. Soc.* **104**, no. 3, pp. 561-574.
- [8] R.L. Dobrushin and P. Major (1979): Non-central limit theorems for nonlinear functionals of Gaussian fields. Z. Wahrscheinlichkeitstheor. Verwandte Geb. 50, pp. 27-52.
- [9] L. Giraitis and D. Surgailis (1985): CLT and other limit theorems for functionals of Gaussian processes. Z. Wahrscheinlichkeitstheor. Verwandte Geb. 70, pp. 191-212.
- [10] K. Gonchigdanzan (2001): Almost Sure Central Limit Theorems. Ph.D. Thesis, University of Cincinnati, Available online.
- [11] I.A. Ibragimov and M.A. Lifshits (1998): On the convergence of generalized moments in almost sure central limit theorem. *Statist. Probab. Lett.* **40**, no. 4, pp. 343-351.
- [12] I.A. Ibragimov and M.A. Lifshits (2000): On limit theorems of "almost sure" type. *Theory Probab. Appl.* 44, no. 2, 254-272.
- [13] M.T. Lacey and W. Philipp (1990): A note on the almost sure central limit theorem. Statist. Probab. Lett. 9, pp. 201-205.
- [14] P. Lévy (1937): Théorie de l'addition des variables aléatoires. Gauthiers-Villars.
- [15] I. Nourdin. Malliavin-Stein approach: a webpage maintained by Ivan Nourdin. http://tinyurl.com/kvpdgcy
- [16] I. Nourdin, D. Nualart and G. Peccati (2016): Strong asymptotic independence on Wiener chaos. *Proc. American Math. Society* **144**, no. 2, pp. 875-886.

- [17] I. Nourdin and G. Peccati (2009): Stein's method on Wiener chaos. *Probab. Theory Related Fields* **145**, no. 1, pp. 75-118.
- [18] I. Nourdin and G. Peccati (2012): Normal Approximations with Malliavin Calculus: From Stein's Method to Universality. Cambridge Tracts in Mathematics. Cambridge University Press.
- [19] I. Nourdin and G. Poly (2013): Convergence in total variation on Wiener chaos. *Stoch. Proc. Appl.* **123**, pp. 651-674.
- [20] I. Nourdin and J. Rosiński (2014): Asymptotic independence of multiple Wiener-Itô integrals and the resulting limit laws. *Ann. Probab.* **42**, no. 2, pp. 497-526.
- [21] D. Nualart and G. Peccati (2005): Central limit theorems for sequences of multiple stochastic integrals. Ann. Probab. 33, no. 1, pp. 177-193.
- [22] D. Nualart (2006): *The Malliavin calculus and related topics*. 2nd edition. Probability and Its Applications, Springer.
- [23] D. Nualart and E. Nualart (2018): Introduction to Malliavin Calculus. Institute of Mathematical Statistics Textbooks, Cambridge University Press.
- [24] M. Peligrad, and Q. M. Shao (1995): A note on the almost sure central limit theorem for weakly dependent random variables. *Statist. Probab. Lett.* **22**, no. 2, pp. 131-136.
- [25] P. Schatte (1988): On strong versions of the central limit theorem. *Math. Nachr.* 137, pp. 249-256.
- [26] F. Spitzer (1976): *Principles of random walk*. Graduate Texts in Mathematics **34**, Second edition, Springer-Verlag, New York-Heidelberg.
- [27] M.S.Taqqu (1979): Convergence of integrated processes of arbitrary Hermite rank. Z. Wahrscheinlichkeitstheor. Verwandte Geb. 50, pp. 53-83.